Table 11.13 All-cause mortality and five main causes of death by smoking status: death rates per 100,000 among men of Cancer Prevention Study II Nutrition Cohort, Health Professional Follow-Up Study, National Household Survey, National Institutes of Health-AARP Diet and Health Study, and Women's Health Initiative, 2000–2010

	Never-smoker			Current smoker			Rate difference (/100,000)	RR (95% CI) current versus never smoker[‡]
	Number of deaths	Person-years	Death rate[†] (/100,000)	Number of deaths	Person-years	Death rate (/100,000)		
All Causes								
Age 55–64	1,182	253,125	401.81	1,170	74,157	1,409.96	1,008.15	2.92 (2.69–3.18)
Age 65–74	6,495	586,441	1,075.35	4,011	110,799	3,619.35	2,544.00	3.00 (2.89–3.13)
Age ≥75	11,312	328,189	4,988.58	1,855	26,789	10,169.60	5,181.03	2.36 (2.24– 2.48)
Lung Cancer[1]								
Age 55–64	34	253,125	12.70	232	74,157	279.94	267.24	19.03 (13.19–27.47)
Age 65–74	147	586,441	24.00	891	110,799	804.05	780.05	28.29 (23.72–33.74)
Age ≥75	190	328,189	69.97	354	26,789	1,557.64	1,487.67	22.52 (18.83–26.92)
COPD[2]								
Age 55–64	2	253,125	0.48	68	74,157	80.61	80.13	84.79 (20.65–348.23)
Age 65–74	55	586,441	9.12	367	110,799	331.13	322.01	29.70 (22.31–39.53)
Age ≥75	137	328,189	61.72	233	26,789	1,378.06	1,316.34	23.01 (18.56–28.51)
Total Stroke[3]								
Age 55–64	40	253,125	13.03	28	74,157	30.52	17.48	2.07 (1.25–3.41)
Age 65–74	310	586,441	50.42	137	110,799	123.61	73.20	2.17 (1.77–2.66)
Age ≥75	748	328,189	356.66	72	26,789	501.14	144.48	1.48 (1.16–1.88)